

# On Egalitarian Belief Merging

**Patricia Everaere**

LIFL-CNRS, Université Lille 1, France  
patricia.everaere@univ-lille1.fr

**Sébastien Konieczny**

CRIL-CNRS, Université d'Artois, France  
{konieczny,marquis}@cril.fr

**Pierre Marquis**

## Abstract

Belief merging aims at defining the beliefs of a group of agents from the beliefs of each member of the group. It is related to more general notions of aggregation from economics (social choice theory). Two main subclasses of belief merging operators exist: majority operators which are related to utilitarianism, and arbitration operators which are related to egalitarianism. Though utilitarian (majority) operators have been extensively studied so far, there is much less work on egalitarian operators. In order to fill the gap, we investigate possible translations in a belief merging framework of some egalitarian properties and concepts coming from social choice theory, such as Sen-Hammond equity, Pigou-Dalton property, median, and Lorenz curves. We study how these properties interact with the standard rationality conditions considered in belief merging. Among other results, we show that the distance-based merging operators satisfying Sen-Hammond equity are mainly those for which *leximax* is used as the aggregation function.

## Introduction

The aim of belief merging is to define a coherent belief base from a set of jointly incoherent belief bases, representing the beliefs of a group of agents. The rationality properties of belief merging operators have been studied in (Revesz 1997; Lin and Mendelzon 1999; Konieczny and Pino Pérez 2002a; Konieczny, Lang, and Marquis 2004; Everaere, Konieczny, and Marquis 2010b). Especially, in (Konieczny and Pino Pérez 2002a), a number of postulates characterizing the so called IC merging operators have been identified. At the same time, many definitions of propositional belief merging operators have been pointed out. Most of these operators are distance-based ones, which means that they can be defined using a distance between interpretations and an aggregation function (Revesz 1997; Lin and Mendelzon 1999; Konieczny, Lang, and Marquis 2004; Everaere, Konieczny, and Marquis 2010a).

Two main subclasses of IC belief merging operators have been defined in (Konieczny and Pino Pérez 2002a): majority

operators, which solve conflicts using majority, and arbitration operators, which try to find a consensual result. However, while many distance-based majority merging operators have been defined in the literature, very few arbitration operators have been identified so far. To be more precise, the only arbitration IC merging operators we are aware of are distance-based operators using *leximax* as aggregation function.

As already discussed in previous merging papers, usual IC merging operators can be used to merge either beliefs or goals. This distinction between belief and goals do not seem to impact any of the usual IC postulates. It is natural to consider egalitarian merging operators when merging goals, if one tries to achieve a “fair” result. This could be important to ensure full adhesion of (autonomous) agents towards the obtained (merged) group goals. As to the belief merging issue, when the aim is to find the correct state of the world, majority methods may seem more appealing. In (Everaere, Konieczny, and Marquis 2010b) we discussed the truth-tracking problem for merging, and we made a distinction between two possible views of belief merging. The one relating to truth-tracking is called the epistemic view, and in this case it is natural to consider only majority operators (this is a consequence of Condorcet Jury Theorem (Everaere, Konieczny, and Marquis 2010b)). But there is a second possible view, called the synthesis view, where the aim is to best represent the opinion of the group (profile). This view has nothing to do with the correct state of the world, the merging process only cares about individual opinions. In this case egalitarian operators prove useful, since they provide in a sense a more robust (i.e. more consensual) view of the opinion of the group than majority operators.

Majority merging operators are closely related to the utilitarian social welfare approaches, where the aim is to determine solutions with the best aggregated utility (Harsanyi 1955; Moulin 1988; Sen 2005). On the other hand, arbitration operators are related to the egalitarian social welfare approaches, where the objective is to find solutions which are as fair as possible; this usually means that they give as much as possible to the poorest agents. To this extent, poverty measures (Rawls 1971; Gini 1921; Sen 1973; Dutta 2002) are relevant to the design of egalitarian approaches.

The aim of this paper is to introduce and study new egal-

itarian merging operators, by exhibiting other fairness conditions than arbitration and by pointing out belief merging operators satisfying them. The methodology we follow to reach our goal consists in investigating equity conditions considered in social choice theory (Arrow, Sen, and Suzumura 2002) in order to determine if they can be reasonably imported in the belief merging setting. Thus, in the following, we translate to the belief merging framework two egalitarian conditions coming from social choice theory: Sen-Hammond equity, and Pigou-Dalton property. We show that the distance-based merging operators satisfying Sen-Hammond equity are mainly those for which *leximax* is used as the aggregation function. We also introduce two new families of belief merging operators, based respectively on the median and on an aggregated sum (Lorenz curves). We identify the rationality properties satisfied by these operators and study in particular their egalitarian behaviour.

The rest of the paper is organized as follows. First we give some preliminaries on propositional belief merging, focusing on IC merging operators and distance-based operators. The egalitarian operators we will define do not satisfy all the usual IC postulates. In fact the usual aggregation postulates **(IC5)** and **(IC6)** are not satisfied. So we define a general family of belief merging operators, called pre-IC merging operators; where these Pareto-related conditions are replaced by unanimity conditions. Then we show how Sen-Hammond equity condition, and Pigou-Dalton property can be expressed in the belief merging setting. Then, we define the family of median distance-based merging operators. We show that the operators of this family based on the *leximed*<sup>k</sup> aggregation functions are pre-IC operators, and those for which  $k \geq 0.5$  satisfy also the arbitration postulate **(Arb)**, but not the Pigou/Dalton property. Finally we introduce the family of cumulative sum distance-based merging operators; we identify in this family some pre-IC operators, and among them an operator satisfying the Pigou-Dalton property.

## On Propositional Belief Merging

We consider a propositional language  $\mathcal{L}$  defined from a finite set of propositional variables  $\mathcal{P}$  and the usual connectives.

An interpretation (or state of the world)  $\omega$  is a total function from  $\mathcal{P}$  to  $\{0, 1\}$ .  $\Omega$  is the set of all interpretations. An interpretation is usually denoted by a bit vector whenever a strict total order on  $\mathcal{P}$  is specified. An interpretation  $\omega$  is a model of a formula  $\phi \in \mathcal{L}$  if and only if it makes it true in the usual truth functional way.  $\models$  and  $\equiv$  denote logical entailment and equivalence, respectively.  $[\phi]$  denotes the set of models of a formula  $\phi$ , i.e.,  $[\phi] = \{\omega \in \Omega \mid \omega \models \phi\}$ .

A base  $K$  denotes the set of beliefs of an agent, it is a finite set of propositional formulae, interpreted conjunctively (i.e., viewed as the conjunction of its elements).

A profile  $E$  denotes a group of  $n$  agents that are involved in the merging process; formally  $E$  is given by a multi-set  $\{K_1, \dots, K_n\}$  of bases.  $\bigwedge E$  denotes the conjunction of all elements of  $E$ , and  $\sqcup$  denotes the multi-set union. Two multi-sets  $E = \{K_1, \dots, K_n\}$  and  $E' = \{K'_1, \dots, K'_n\}$  are equivalent, noted  $E \equiv E'$ , iff there exists a permutation  $\pi$  over  $\{1, \dots, n\}$  s.t. for each  $i \in 1, \dots, n$ , we have

$$K_i \equiv K'_{\pi(i)}.$$

An *integrity constraint*  $\mu$  is a formula restricting the possible results of the merging process.

A *merging operator*  $\Delta$  is a function which associates with a profile  $E$  and an integrity constraint  $\mu$  a merged base  $\Delta_\mu(E)$ .

The logical properties given in (Konieczny and Pino Pérez 2002a) for characterizing IC belief merging operators are:

**Definition 1** A merging operator  $\Delta$  is an IC merging operator iff it satisfies the following properties:

- (IC0)**  $\Delta_\mu(E) \models \mu$
- (IC1)** If  $\mu \not\models \perp$ , then  $\Delta_\mu(E) \not\models \perp$
- (IC2)** If  $\bigwedge E \wedge \mu \not\models \perp$ , then  $\Delta_\mu(E) \equiv \bigwedge E \wedge \mu$
- (IC3)** If  $E_1 \equiv E_2$  and  $\mu_1 \equiv \mu_2$ , then  $\Delta_{\mu_1}(E_1) \equiv \Delta_{\mu_2}(E_2)$
- (IC4)** If  $K_1 \models \mu$  and  $K_2 \models \mu$ , then  $\Delta_\mu(\{K_1, K_2\}) \wedge K_1 \not\models \perp$  if and only if  $\Delta_\mu(\{K_1, K_2\}) \wedge K_2 \not\models \perp$
- (IC5)**  $\Delta_\mu(E_1) \wedge \Delta_\mu(E_2) \models \Delta_\mu(E_1 \sqcup E_2)$
- (IC6)** If  $\Delta_\mu(E_1) \wedge \Delta_\mu(E_2) \not\models \perp$ , then  $\Delta_\mu(E_1 \sqcup E_2) \models \Delta_\mu(E_1) \wedge \Delta_\mu(E_2)$
- (IC7)**  $\Delta_{\mu_1}(E) \wedge \mu_2 \models \Delta_{\mu_1 \wedge \mu_2}(E)$
- (IC8)** If  $\Delta_{\mu_1}(E) \wedge \mu_2 \not\models \perp$ , then  $\Delta_{\mu_1 \wedge \mu_2}(E) \models \Delta_{\mu_1}(E)$

See (Konieczny and Pino Pérez 2002a) for explanations on these properties. Two subclasses of IC merging operators are also defined in (Konieczny and Pino Pérez 2002a):

**Definition 2** An IC majority operator is an IC merging operator which satisfies the following majority property:

$$\mathbf{(Maj)} \quad \exists n \quad \Delta_\mu(E_1 \sqcup \underbrace{E_2 \sqcup \dots \sqcup E_2}_n) \models \Delta_\mu(E_2)$$

An IC arbitration operator is an IC merging operator which satisfies the following arbitration property:<sup>1</sup>

$$\mathbf{(Arb)} \quad \text{If } \begin{cases} \Delta_{\mu_1}(K_1) \equiv \Delta_{\mu_2}(K_2) \\ \Delta_{\mu_1 \leftrightarrow \neg \mu_2}(\{K_1, K_2\}) \equiv (\mu_1 \leftrightarrow \neg \mu_2) \\ \mu_1 \not\models \mu_2 \\ \mu_2 \not\models \mu_1 \end{cases} \text{ then } \Delta_{\mu_1 \vee \mu_2}(\{K_1, K_2\}) \equiv \Delta_{\mu_1}(K_1)$$

Majority operators solve conflicts using majority. **(Maj)** says that if one duplicates sufficiently many times a profile  $E_2$ , then the result of the merging of  $E_1$  with the  $E_2$  duplications will obey the choices of the profile  $E_2$ . Arbitration operators try to find a consensual result. See the corresponding condition 8 in Definition 3 and Example 1 which illustrates how this property gives a preference to consensual (“median”) choices of interpretations.

The representation theorems enable to interpret these logical properties as constraints on the choice of interpretations for defining the models of the resulting base:

**Definition 3** A syncretic assignment is a function mapping each profile  $E$  to a total pre-order  $\leq_{E^2}$  over  $\Omega$  such that for

<sup>1</sup>When  $E = \{K\}$  we note  $\Delta_\mu(K)$  instead of  $\Delta_\mu(\{K\})$ .

<sup>2</sup>For every pre-order  $\leq$ ,  $<$  denotes its strict part and  $\simeq$  the corresponding indifference relation. Furthermore, we will use  $\leq_K$  as a short for  $\leq_{\{K\}}$ , and  $\leq_\omega$  as a short for any  $\leq_{\{K_\omega\}}$  where  $\omega$  is the unique model of  $K_\omega$ .

any profiles  $E, E_1, E_2$  and for any bases  $K, K'$  the following conditions hold:

1. If  $\omega \models \bigwedge E$  and  $\omega' \models \bigwedge E$ , then  $\omega \simeq_E \omega'$
2. If  $\omega \models \bigwedge E$  and  $\omega' \not\models \bigwedge E$ , then  $\omega <_E \omega'$
3. If  $E_1 \equiv E_2$ , then  $\leq_{E_1} = \leq_{E_2}$
4.  $\forall \omega \models K \exists \omega' \models K' \omega' \leq_{\{K, K'\}} \omega$
5. If  $\omega \leq_{E_1} \omega'$  and  $\omega \leq_{E_2} \omega'$ , then  $\omega \leq_{E_1 \sqcup E_2} \omega'$
6. If  $\omega <_{E_1} \omega'$  and  $\omega \leq_{E_2} \omega'$ , then  $\omega <_{E_1 \sqcup E_2} \omega'$

A majority syncretic assignment is a syncretic assignment which satisfies the following condition:

7. If  $\omega <_{E_2} \omega'$ , then  $\exists n \omega <_{E_1 \sqcup E_2^n} \omega'$

A fair syncretic assignment is a syncretic assignment which satisfies the following condition:

8. If  $\omega <_{K_1} \omega'$ ,  $\omega <_{K_2} \omega''$ , and  $\omega' \simeq_{\{K_1, K_2\}} \omega''$ , then  $\omega <_{\{K_1, K_2\}} \omega''$

**Proposition 1 ((Konieczny and Pino Pérez 2002a))** A merging operator  $\Delta$  is an IC merging operator (resp. an IC majority, an IC arbitration operator) iff there exists a syncretic assignment (resp. a majority syncretic assignment, a fair syncretic assignment) that maps each profile  $E$  to a total pre-order  $\leq_E$  over  $\Omega$  such that  $[\Delta_\mu(E)] = \min([\mu], \leq_E)$ .

Let us now give some examples of IC merging operators using the family of distance-based merging operators (Konieczny, Lang, and Marquis 2004):

**Definition 4** A distance<sup>3</sup> between interpretations is a function  $d : \Omega \times \Omega \rightarrow \mathbb{R}^+$  such that for any  $\omega_1, \omega_2 \in \Omega$ :

- $d(\omega_1, \omega_2) = d(\omega_2, \omega_1)$
- $d(\omega_1, \omega_2) = 0$  iff  $\omega_1 = \omega_2$

Usual distances considered in merging are the Hamming distance  $d_H$ :  $d_H(\omega_1, \omega_2)$  is the number of propositional letters on which the two interpretations differ (this corresponds to the 1-norm distance, also referred to as the Manhattan distance) and the drastic distance  $d_D$ , defined as  $d_D(\omega_1, \omega_2) = 0$  if  $\omega_1 = \omega_2$ , and  $= 1$  otherwise (this corresponds to the infinity-norm distance, also known as Chebyshev distance).

**Definition 5** An aggregation function is a mapping<sup>4</sup>  $f$  from  $\mathbb{R}^m$  to  $\mathbb{R}$ , which satisfies:

- if  $x_i \geq x'_i$ , then  $f(x_1, \dots, x_i, \dots, x_m) \geq f(x_1, \dots, x'_i, \dots, x_m)$  **(non-decreasingness)**
- $f(x_1, \dots, x_m) = 0$  if  $\forall i, x_i = 0$  **(minimality)**
- $f(x) = x$  **(identity)**
- If  $\sigma$  is a permutation over  $\{1, \dots, m\}$ , then  $f(x_1, \dots, x_m) = f(x_{\sigma(1)}, \dots, x_{\sigma(m)})$  **(symmetry)**

Some additional properties can be considered for  $f$ , especially:

- if  $x_i > x'_i$ , then  $f(x_1, \dots, x_i, \dots, x_m) > f(x_1, \dots, x'_i, \dots, x_m)$  **(strict non-decreasingness)**

<sup>3</sup>More exactly pseudo-distances since triangular inequality is not required, but for the sake of simplicity we will abuse words in this way.

<sup>4</sup>Strictly speaking, it is a family of mappings, one for each  $m$ .

**Definition 6** Let  $d$  and  $f$  be respectively a distance between interpretations and an aggregation function. The distance-based merging operator  $\Delta^{d,f}$  is defined by  $[\Delta_\mu^{d,f}(E)] = \min([\mu], \leq_E)$ , where the total pre-order  $\leq_E$  on  $\Omega$  is defined in the following way (with  $E = \{K_1, \dots, K_n\}$ ):

- $\omega \leq_E \omega'$  iff  $d(\omega, E) \leq d(\omega', E)$
- $d(\omega, E) = f(d(\omega, K_1), \dots, d(\omega, K_n))$
- $d(\omega, K) = \min_{\omega' \models K} d(\omega, \omega')$

For usual aggregation functions, whatever the chosen distance, the corresponding distance-based operators exhibit good logical properties ((Konieczny and Pino Pérez 2002b; Everaere, Konieczny, and Marquis 2010a)):

**Proposition 2** For any distance  $d$ :

- if  $f$  is the sum  $\Sigma$ , *leximin*<sup>5</sup>, or  $\Sigma^n$  (the sum of the  $n^{\text{th}}$  powers), then  $\Delta^{d,f}$  is an IC majority operator.
- if  $f$  is *leximax*, then  $\Delta^{d,f}$  is an IC arbitration operator.

## Pre-IC Operators

We now define a general family of belief merging operators, called pre-IC merging operators, obtained by relaxing the two postulates **(IC5)** and **(IC6)** into two natural conditions used in other aggregation theories contexts.

**Definition 7** A merging operator  $\Delta$  is pre-IC merging operator iff it satisfies **(IC0)** to **(IC4)**, **(IC7)** to **(IC8)** and the following properties:

**(IC5b)**  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n) \models \Delta_\mu(\{K_1, \dots, K_n\})$

**(IC6b)** If  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n) \not\models \perp$ , then  $\Delta_\mu(\{K_1, \dots, K_n\}) \models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$

Thus, switching from IC operators to pre-IC ones simply consists in replacing the postulates **(IC5)** and **(IC6)** postulates by the weaker postulates **(IC5b)** and **(IC6b)**. Indeed, it is easy to prove that **(IC5b)** (resp. **(IC6b)**) is implied by **(IC5)** (resp. **(IC6)**). As a consequence, we have:

**Proposition 3** Every IC merging operator is a pre-IC merging operator.

Let us now give a representation theorem suited to the pre-IC family:

**Definition 8** A pre-syncretic assignment is a function mapping each profile  $E$  to a total pre-order  $\leq_E$  over  $\Omega$  such that for any profiles  $E, E_1, E_2$  and for any bases  $K, K'$ , the following conditions hold:

1. If  $\omega \models E$  and  $\omega' \models E$  then  $\omega \equiv_E \omega'$
2. If  $\omega \models E$  and  $\omega' \not\models E$  then  $\omega <_E \omega'$
3. If  $E_1 \equiv E_2$  then  $\leq_{E_1} = \leq_{E_2}$
4.  $\forall \omega \models K, \exists \omega' \models K' \omega' \leq_{\{K, K'\}} \omega$
- 5b. If  $\forall i \omega \leq_{K_i} \omega'$  then  $\omega \leq_{\{K_1, \dots, K_n\}} \omega'$
- 6b. If  $\forall i \omega \leq_{K_i} \omega'$  and  $\exists k \omega <_{K_k} \omega'$  then  $\omega <_{\{K_1, \dots, K_n\}} \omega'$

<sup>5</sup>The *leximin* (resp. *leximax*) aggregation function selects the interpretations that are minimal for the lexicographic order, once the distances are sorted into the increasing (resp. decreasing) order.

Conditions 5b and 6b are direct translations of Pareto conditions, which are usual conditions in social choice, multi-criteria decision making, etc. So they should be considered as minimal aggregation conditions to be satisfied. Conditions 5 and 6 (Definition 3) are much more demanding, since they constraint all unions of two profiles.

**Proposition 4** *A merging operator  $\Delta$  is a pre-IC merging operator iff there exists a pre-syncretic assignment that maps each profile  $E$  to a total pre-order  $\leq_E$  over  $\Omega$  such that*

$$[\Delta_\mu(E)] = \min([\mu], \leq_E).$$

**Proof: Only If** Let  $\Delta$  be a pre-IC merging operator. We associate with it a pre-syncretic assignment as follows: for any profile  $E = \{K_1, \dots, K_n\}$ , we define  $\leq_E$  by  $\omega \leq_E \omega'$  if and only if  $\omega \models \Delta_\mu(E)$  where  $[\mu] = \{\omega, \omega'\}$ . From (Konieczny and Pino Pérez 2002a), we know that  $\leq_E$  is a total pre-order; furthermore, as  $\Delta$  satisfies (IC0-IC4) and (IC7-IC8), conditions 1 to 4 are satisfied by the assignment. We have then only to check that conditions 5b and 6b are satisfied. Consider two interpretations  $\omega$  and  $\omega'$ , and let  $\mu$  be such that  $[\mu] = \{\omega, \omega'\}$ :

Condition 5b: suppose that  $\forall i \in \{1, \dots, n\}, \omega \leq_{K_i} \omega'$ . Then  $\forall i \in \{1, \dots, n\}, \omega \models \Delta_\mu(K_i)$ . Stated otherwise,  $\omega \models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . From (IC5b), we know that  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n) \models \Delta_\mu(\{K_1, \dots, K_n\})$ , so  $\omega \models \Delta_\mu(\{K_1, \dots, K_n\})$ . Therefore,  $\omega \leq_E \omega'$ , hence condition 5b is satisfied.

Condition 6b: suppose that  $\forall i \in \{1, \dots, n\}, \omega \leq_{K_i} \omega'$  and  $\exists k \in \{1, \dots, n\}, \omega <_{K_k} \omega'$ . Then  $\forall i \in \{1, \dots, n\}, \omega \models \Delta_\mu(K_i)$  and  $\omega' \not\models \Delta_\mu(K_k)$ . Thus we have  $\omega \models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$  and  $\omega' \not\models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . Hence  $[\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)] = \{\omega\}$ . From (IC6b), since  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$  is consistent, it must be the case that  $\Delta_\mu(\{K_1, \dots, K_n\}) \models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . Since  $\mu$  is consistent and  $\Delta$  satisfies (IC1),  $\Delta_\mu(\{K_1, \dots, K_n\})$  must be consistent as well. As a consequence, we must have  $[\Delta_\mu(\{K_1, \dots, K_n\})] = \{\omega\}$ . Therefore,  $\omega \models \Delta_\mu(\{K_1, \dots, K_n\})$  and  $\omega' \not\models \Delta_\mu(\{K_1, \dots, K_n\})$ , showing that  $\omega <_E \omega'$ . Hence condition 6b is satisfied.

**If** Consider a pre-syncretic assignment mapping each belief set  $E$  to a total pre-order  $\leq_E$  over interpretations. Consider the merging operator  $\Delta$  defined as follows:  $[\Delta_\mu(E)] = \min([\mu], \leq_E)$ . From (Konieczny and Pino Pérez 2002a), we know that  $\Delta$  satisfies (IC0-IC4) and (IC7-IC8). Let us now show that  $\Delta$  satisfies (IC5b-IC6b). Suppose that  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$  is consistent (if not, (IC5b-IC6b) is trivially satisfied).

**(IC5b)** let  $\omega$  be any model of  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . So  $\forall \omega' \models \mu, \omega \leq_{K_i} \omega'$ . From condition 5b, we know that  $\forall \omega' \models \mu, \omega \leq_{\{K_1, \dots, K_n\}} \omega'$ . Hence by definition of  $\Delta$ , we have  $\omega \models \Delta_\mu(\{K_1, \dots, K_n\})$ . Since this holds for any model  $\omega$  of  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ , we conclude that  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n) \models \Delta_\mu(\{K_1, \dots, K_n\})$  and (IC5b) is satisfied.

**(IC6b)** towards a contradiction, suppose that  $\Delta_\mu(\{K_1, \dots, K_n\}) \not\models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . So  $\exists \omega$  s.t.  $\omega \models \Delta_\mu(\{K_1, \dots, K_n\})$  and  $\omega \not\models \Delta_\mu(K_1)$

$\wedge \dots \wedge \Delta_\mu(K_n)$ . Since  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$  is consistent, there is an interpretation  $\omega'$  which is a model of  $\Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ . Hence  $\forall i \in \{1, \dots, n\}, \omega' \leq_{K_i} \omega$ . Since  $\omega \not\models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ ,  $\exists k \in \{1, \dots, n\}$  s.t.  $\omega \not\models \Delta_\mu(K_k)$ , showing that  $\omega' <_{K_k} \omega$ . From condition 6b, we conclude that  $\omega' <_{\{K_1, \dots, K_n\}} \omega$ . This contradicts the fact that  $\omega \models \Delta_\mu(\{K_1, \dots, K_n\})$ . Hence,  $\Delta_\mu(\{K_1, \dots, K_n\}) \models \Delta_\mu(K_1) \wedge \dots \wedge \Delta_\mu(K_n)$ , and (IC6b) is satisfied.  $\square$

As one can expect, it is much easier to satisfy the pre-IC merging conditions than IC merging ones. We can for instance show that:

**Proposition 5** *If  $d$  is any distance and  $f$  is any aggregation function satisfying strict non-decreasingness, then the merging operator  $\Delta^{d,f}$  is a pre-IC merging operator.*

One can compare this result with a corresponding one about distance-based IC merging operators, reported in (Konieczny, Lang, and Marquis 2004), and showing that the aggregation function  $f$  has to satisfy two additional conditions in order to guarantee that the distance-based merging operator given by  $d$  and  $f$  is an IC merging operator.

## Conditions for Egalitarian Merging

The only egalitarian property that has been proposed so far for belief merging is the *arbitration* property, represented by the **(Arb)** postulate (or the corresponding semantic condition 8 on syncretic assignments). So a key issue we would like to address is to determine whether other egalitarian properties are possible in the belief merging framework, and, if so, how they relate with arbitration.

If one looks closely at condition 8 in Definition 3, it is clear that the arbitration property only imposes some constraints on the merging of profiles consisting of two bases. Then the IC properties **(IC5)** and **(IC6)** ensure the propagation of the constraints on profiles of any size.

We now propose a first alternative condition, coming from social choice theory, for characterizing egalitarian behaviour in belief merging. This condition, proposed by Hammond (1976) is known in the literature as the Sen-Hammond equity condition (Sen 1997; Suzumura 1983). This condition can be expressed in the following way:

**Definition 9 (Sen97)** *If person  $i$  is worse off than person  $j$  both in  $x$  and in  $y$ , and if  $i$  is better off himself in  $x$  than in  $y$ , while  $j$  is better off in  $y$  than in  $x$ , and if furthermore all others are just as well off in  $x$  as in  $y$ , then  $x$  is socially at least as good as  $y$ .*

It is translated in the belief merging setting as constraints on the total pre-orders associated with the input profiles. These constraints concern profiles of arbitrary size, and not only those consisting of two bases. Before doing it, we first need to define a notion of the respective ‘‘satisfaction’’ of two bases given an interpretation:

**Definition 10** *Given a merging operator  $\Delta$  defining an assignment which maps every profile  $E$  to a total pre-order  $\leq_E$  over  $\Omega$ , given an interpretation  $\omega$ , and two bases  $K_1$*

and  $K_2$ , we say that  $K_1$  is better than  $K_2$  given  $\omega$ , denoted  $K_1 \prec_\omega K_2$ , iff  $\exists \omega_1 \models K_1, \forall \omega_2 \models K_2, \omega_1 <_\omega \omega_2$ .

We can now give the translation of the Sen-Hammond Equity (SHe) condition:

**Definition 11 (Condition (SHe))** Let  $E = \{K_1, \dots, K_n\}$ .

$$\left. \begin{array}{l} \omega <_{K_1} \omega' \\ \omega' <_{K_2} \omega \\ \forall i \neq 1, 2 \omega \simeq_{K_i} \omega' \\ K_1 \prec_\omega K_2 \\ K_1 \prec_{\omega'} K_2 \end{array} \right\} \implies \omega' \leq_E \omega$$

When distance-based merging operators are considered, this condition is equivalent to:

**Definition 12 (Condition (SHE))**

$$\left. \begin{array}{l} d(\omega, K_1) < d(\omega', K_1) < d(\omega', K_2) < d(\omega, K_2) \\ \forall i \neq 1, 2 d(\omega, K_i) = d(\omega', K_i) \end{array} \right\} \implies$$

$$f(d(\omega', K_1), d(\omega', K_2), \dots, d(\omega', K_n)) \leq f(d(\omega, K_1), d(\omega, K_2), \dots, d(\omega, K_n))$$

**Proposition 6** A distance-based merging operator  $\Delta^{d,f}$  satisfies (SHE) if and only if it satisfies (SHe).

**Proof:** We prove the equivalence between (SHe) and (SHE) by showing that those two implications have equivalent premises and equivalent conclusions.

By definition  $K_1 \prec_\omega K_2$  iff  $\exists \omega_1 \models K_1, \forall \omega_2 \models K_2, \omega_1 <_\omega \omega_2$ . Then  $\exists \omega_1 \models K_1, \forall \omega_2 \models K_2, d(\omega, \omega_1) < d(\omega, \omega_2)$ :  $\exists \omega_1 \models K_1, d(\omega, \omega_1) < \min_{\omega_2 \models K_2} d(\omega, \omega_2) = d(\omega, K_2)$ . Hence we have  $d(\omega, K_1) < d(\omega, K_2)$ .

$$\begin{array}{l} \omega <_{K_1} \omega' \Leftrightarrow d(\omega, K_1) < d(\omega', K_1) \\ \omega' <_{K_2} \omega \Leftrightarrow d(\omega', K_2) < d(\omega, K_2) \\ \forall i \neq 1, 2 \omega \simeq_{K_i} \omega' \Leftrightarrow \forall i \neq 1, 2 d(\omega, K_i) = d(\omega', K_i) \\ K_1 \prec_\omega K_2 \Leftrightarrow d(\omega, K_1) < d(\omega, K_2) \\ K_1 \prec_{\omega'} K_2 \Leftrightarrow d(\omega', K_1) < d(\omega', K_2) \end{array}$$

This shows the equivalence of the premises. Now, using transitivity, we get:  $d(\omega, K_1) < d(\omega', K_1) < d(\omega', K_2) < d(\omega, K_2)$  and  $\forall i \neq 1, 2 \omega \simeq_{K_i} \omega' \Leftrightarrow \forall i \neq 1, 2 d(\omega, K_i) = d(\omega', K_i)$ . Hence  $\omega' \leq_E \omega \Leftrightarrow f(d(\omega', K_1), d(\omega', K_2), \dots, d(\omega', K_n)) \leq f(d(\omega, K_1), d(\omega, K_2), \dots, d(\omega, K_n))$ .  $\square$

This Sen-Hammond Equity condition expresses the following idea: compare two ‘‘situations’’  $\omega_1$  and  $\omega_2$  that are equally good for all the agents except two of them. For these two agents one of them ( $K_1$ ) is in the two situations better than the other agent ( $K_2$ ). Then the fairer situation is the one that gives the more to the less satisfied agent ( $K_2$ ).

This condition looks close to the arbitration condition (compare (SHe) with condition 8 of Definition 3), but the two conditions are logically independent. Let us now illustrate on a simple example how they differ:

**Example 1** Consider a propositional language over two variables and a distance-based merging operator  $\Delta^{d,f}$ . Suppose that the distance  $d$  on which  $\Delta^{d,f}$  is built is the shortest path distance on the following graph:



Figure 1: Egalitarian behaviour - (Arb)




Figure 2: Egalitarian behaviour - (SHE)

$$\omega_1 \xrightarrow{1} \omega_2 \xrightarrow{1} \omega_3 \xrightarrow{1} \omega_4.$$

Figure 1 illustrates the case when the unique model of  $K_1$  is  $\omega_1$  and the unique model of  $K_2$  is  $\omega_3$ . It is clear here that  $\omega_1$  and  $\omega_3$  play symmetrical roles, so that they cannot be distinguished when  $K_1$  and  $K_2$  are merged. However, on this example,  $\omega_2$  appears as a more consensual choice than  $\omega_1$  and  $\omega_3$  for this merging. Accordingly, (Arb) imposes that  $\omega_2$  is strictly preferred to  $\omega_1$  and to  $\omega_3$  w.r.t.  $\leq_{\{K_1, K_2\}}$ .

Suppose now that we want to merge  $E' = \{K'_1, K'_2\}$  where  $[K'_1] = \{\omega_1, \omega_3\}$  and  $[K'_2] = \{\omega_4\}$ , and that the choice to be made is between  $\omega_1$  and  $\omega_2$  (i.e., the integrity constraint  $\mu$  satisfies  $[\mu] = \{\omega_1, \omega_2\}$ ). This is illustrated by Figure 2. (Arb) imposes no constraint on this choice. But clearly, whatever the choice (between  $\{\omega_1\}$ ,  $\{\omega_2\}$ , or  $\{\omega_1, \omega_2\}$ ), the result will be closer to  $K'_1$  than to  $K'_2$ . So a simple equity argument is to commit to the choice that is the best for the furthest base  $K'_2$ , so to consider that  $\omega_2$  is strictly preferred to  $\omega_1$  in  $\leq_{E'}$ . This is what (SHe) gives.

Unfortunately, the family of distance-based IC operators satisfying condition (SHE) looks rather limited:

**Proposition 7** Let  $d$  be any distance and  $f$  be any aggregation function satisfying strict non-decreasingness. The IC merging operator  $\Delta^{d,f}$  satisfies condition (SHE) if and only if  $f = \text{leximax}$ .

**Proof:** We know that  $\Delta^{d, \text{leximax}}$  is an IC operator and that  $\text{leximax}$  satisfies strict-decreasingness. It is easy to check that  $\Delta^{d, \text{leximax}}$  satisfies (SHE), so we have mainly to prove the converse implication.

Consider a distance  $d$  and an aggregation function  $f$  satisfying strict non-decreasingness, and suppose that  $\Delta^{d,f}$  satisfies (SHE).

Let  $X = (x_1, \dots, x_n)$  and  $X' = (x'_1, \dots, x'_n)$  be two vectors of distances to a profile  $E$ , ordered in the descending way ( $\forall i, x_i = d(\omega, K_i)$  and  $x_i \geq x_{i+1}$ ;  $x'_i = d(\omega', K_i)$  and  $x'_i \geq x'_{i+1}$ ). We have to show that:

$$X <_{\text{leximax}} X' \Leftrightarrow f(X) < f(X') \quad (1)$$

and

$$X \simeq_{leximax} X'^6 \Leftrightarrow f(X) = f(X') \quad (2)$$

We start with statement (1). Suppose that  $X <_{leximax} X'$ . We know that  $\exists l$  s.t.  $\forall i < l, x_i = x'_i$  and  $x_l < x'_l$ .

**Case 1:**  $l = n$  or  $\forall i > l, x_i = x'_i$ : using strict non-decreasingness we get  $f(X) < f(X')$ .

**Case 2:** Suppose that  $\exists k > l, x_k \neq x'_k$  and  $\forall i \neq k, l, x_i = x'_i$ .

- If  $x_k \leq x'_k$ , then using strict-decreasingness (as  $x_l < x'_l$ ), we get  $f(X) < f(X')$ .
- If  $x_k > x'_k$ , then we have  $x'_l > x_l \geq x_k > x'_k$ . Consider a vector  $Y = (y_1, \dots, y_n)$  s.t.  $\forall i \neq k, l, y_i = x'_i$  and  $y_k, y_l$  such that  $x'_l > y_l > y_k > x_l$ . As we have  $\forall i \neq k, l, y_i = x'_i$  and  $x'_l > y_l > y_k > x'_k$ , because  $\Delta^{d,f}$  satisfies **(SHE)**, we can conclude that  $f(Y) \leq f(X')$ . Furthermore  $\forall i \neq k, l, y_i = x_i$  and  $y_l > y_k > x_l \geq x_k$ , using strict decreasingness, we get  $f(Y) > f(X)$ . By transitivity, we obtain  $f(X) < f(X')$ .

**Case 3:** Let us consider now the general case.  $\forall i < l, x_i = x'_i, x_l < x'_l$  and  $l < n$ . We define  $n - l + 1$  vectors  $Y^r = (y_1^r, \dots, y_n^r)$ , for  $r = 0$  to  $n - l$ :

$$y_i^r = \begin{cases} x_i & \text{if } i < l \\ x_l + \frac{r}{n-l}(x'_l - x_l) & \text{if } i = l \\ x_i & \text{if } l + 1 \leq i \leq n - r \\ x'_i & \text{if } n - r + 1 \leq i \leq n \end{cases}$$

We have  $Y^0 = X$  and  $Y^{n-l} = X'$ , and any two consecutive vectors  $Y^r$  and  $Y^{r+1}$  differ only on two components, namely  $y_l^r$  and  $y_l^{r+1}$  on the one hand, and  $y_{n-r}^r$  and  $y_{n-r}^{r+1}$  on the other hand. Furthermore,  $Y^r <_{leximax} Y^{r+1}$ , because  $\forall i < l, y_i^r = y_i^{r+1}, \forall i \leq l, y_i^r \geq y_i^{r+1}, \forall i \leq l, y_i^{r+1} \geq y_i^{r+1}$  and  $y_l^r < y_l^{r+1}$ . Using Case 2, we can conclude that  $f(Y^r) < f(Y^{r+1})$ . By transitivity we get  $f(Y^0) = f(X) < f(Y^{n-l}) = f(X')$ , and the conclusion follows.

Suppose now that  $f(X) < f(X')$ . If  $X \simeq_{leximax} X'$ , then  $X = X'$  and  $f(X) = f(X')$ : contradiction. If  $X >_{leximax} X'$ , then taking advantage of the first part of the proof, we get  $f(X) > f(X')$ : contradiction. So  $X <_{leximax} X'$ .

Consider now statement (2).

If  $X \simeq_{leximax} X'$ , then  $X = X'$  and  $f(X) = f(X')$ .

Suppose now  $f(X) = f(X')$ . If  $X >_{leximax} X'$ , then taking advantage of the first part of the proof, we get  $f(X) > f(X')$ : contradiction. If  $X <_{leximax} X'$ , then taking advantage of the first part of the proof, we get  $f(X) < f(X')$ : contradiction. So  $X \simeq_{leximax} X'$ .  $\square$

Let us stress here that the condition of strict non-decreasingness is quite natural and not very demanding. Actually all the aggregation functions giving rise to IC merging

<sup>6</sup>If  $X \simeq_{leximax} X'$ , then  $\forall i, x_i = x'_i$ , so  $X = X'$ .

operators we are aware of (including  $\Sigma$ , *leximax*, *leximin*,  $\Sigma^n$ , etc.) satisfy non-decreasingness.

Given this proposition, defining other egalitarian distance-based merging operators requires to focus on other equity principles, or to weaken some IC postulates. We explore both ways in the following.

Thus, we first focus on another egalitarian condition from the social choice literature, namely Pigou-Dalton transfer principle (Dalton 1920). The idea underlying it is that every transfer from the most satisfied agent to the least satisfied one decreases the inequalities:

**Definition 13** *Let  $f$  be an aggregation function.  $f$  satisfies the Pigou-Dalton condition if for all vectors  $X = (x_1, \dots, x_n)$  and  $X' = (x'_1, \dots, x'_n)$ , if  $x_1 < x'_1 \leq x'_2 < x_2$  and  $x'_1 - x_1 = x_2 - x'_2$  and  $\forall i \neq 1, 2, x_i = x'_i$  then  $f(X) < f(X')$ .*

This principle states that if  $X'$  can be obtained from  $X$  by just making some satisfaction transfer from a well-satisfied agent to a less satisfied one, without changing the fact that the first one is still more satisfied than the second one, then  $X'$  is fairer than  $X$ .

This principle can be translated as follows for distance-based merging:

**Definition 14 (Condition (PD))** *If  $\exists k$  and  $l$  s.t.  $d(\omega, K_k) < d(\omega', K_k) \leq d(\omega', K_l) < d(\omega, K_l)$  and  $d(\omega', K_k) - d(\omega, K_k) = d(\omega, K_l) - d(\omega', K_l)$  and  $\forall i \neq k$  and  $i \neq l, d(\omega, K_i) = d(\omega', K_i)$  then  $\omega' <_E \omega$ .*

Of course not all distance-based IC operators satisfy the **(PD)** condition. However, it is satisfied by the well-known arbitration operators based on *leximax*:

**Proposition 8** *Let  $d$  be any distance.*

- $\Delta^{d,leximax}$  satisfies the **(PD)** condition.
- $\Delta^{d,\Sigma}$  and  $\Delta^{d,leximin}$  do not satisfy the **(PD)** condition.

## Median Operators

In this section we define a new family of merging operators using generalized median aggregation functions. Interestingly, some operators of this family are pre-IC merging operators and they satisfy **(Arb)**. The idea of using the median value is very motivated by trying to be as fair as possible. Instead of focusing on a unique aggregation function, we study a full family of  $k$ -median aggregation functions, inspired by the phantom voters voting rules of (Moulin 1988).

**Definition 15** *Let  $k \in ]0, 1]$  be a real number, the  $k$ -median  $med^k(\{x_1, \dots, x_n\})$  of a multi-set  $X = \{x_1, \dots, x_n\}$  of values from a totally ordered set, is the value  $m^k = x_{\sigma(\lceil n \cdot k \rceil)}$  of  $X$ , where  $\sigma$  is a permutation of  $X$  where the  $x_i$  are sorted in ascending order ( $\lceil \cdot \rceil$  denotes the ceiling function).*

For  $k = 0.5$ , the usual notion of median is retrieved.

In many cases these  $k$ -median functions  $med^k$  are not discriminative enough, just like *min* and *max* functions. So, we can define  $k$ -leximedians, noted *k-leximed*, which are to  $med^k$  what *leximin* (resp. *leximax*) is to *min* (resp. *max*).

	$K_1$	$K_2$	$K_3$	distance vector	$med^{0.5}$
000	0	1	1	(0, 1, 1)	1
001	1	0	2	(0, 1, 2)	1
010	1	2	2	(1, 2, 2)	2
011	2	1	3	(1, 2, 3)	2
<b>100</b>	<b>0</b>	<b>2</b>	<b>0</b>	<b>(0, 0, 2)</b>	<b>0</b>
101	1	1	1	(1, 1, 1)	1
110	1	3	1	(1, 1, 3)	1
111	2	2	2	(2, 2, 2)	2

Table 1: Merging with  $\Delta^{d_H, med^{0.5}}$

**Definition 16** Let  $L_1$  and  $L_2$  be two multi-sets consisting of  $n$  elements from a totally ordered set:

$L_1 \leq_{leximed}^k L_2$  iff

- $med^k(L_1) < med^k(L_2)$  or
- $med^k(L_1) = med^k(L_2)$  and  $L_1 \setminus \{med^k(L_1)\} \leq_{leximed}^k L_2 \setminus \{med^k(L_2)\}$

Let us define the  $k$ -median merging operators, and later the  $k$ -leximedians:

**Definition 17** Let  $E = \{K_1, \dots, K_n\}$  be a profile,  $d$  a distance between interpretations and  $k \in ]0, 1]$  a real number. Let  $d_{med}^{d,k}(\omega, E) = med^k(d(\omega, K_1), d(\omega, K_2), \dots, d(\omega, K_n))$ . We define  $[\Delta_{\mu}^{d, med^k}(E)]$  by

$$[\Delta_{\mu}^{d, med^k}(E)] = \{\omega \models \mu \mid d_{med}^k(\omega, E) \text{ is minimal}\}.$$

Here is an example illustrating the behaviour of  $k$ -median operators:

**Example 2** We consider a profile  $E$  of three bases such that  $[K_1] = \{000, 100\}$ ,  $[K_2] = \{001\}$  and  $[K_3] = \{100\}$ . There is no integrity constraint ( $\mu = \top$ ), we use the Hamming distance  $d_H$  and the value  $k = 0.5$  for the “standard” median. The computations are presented in Table 1.

We get  $[\Delta_{\top}^{d_H, med^{0.5}}(E)] = \{100\}$ . The only selected interpretation is 100, because the best vector is (0, 0, 2), with a median value of 0.

Some specific values of  $k$  lead to well-known merging operators: given a profile  $E$  of  $n$  bases,  $\Delta^{med^{\epsilon}}$ , with  $\epsilon \in ]0, \frac{1}{n}]$  corresponds to the *min* operator, and  $\Delta^{med^{\alpha}}$ , with  $\alpha \in ]\frac{n-1}{n}, 1]$  to the *max* operator.

Let us now make precise the rationality postulates satisfied by these operators:

**Proposition 9** For any real number  $k \in ]0, 1]$  and any distance  $d$ ,  $\Delta^{d, med^k}$  satisfies **(IC0)**, **(IC1)**, **(IC3)**, **(IC4)**, **(IC7)**, **(IC8)** and **(IC5b)**. **(IC2)**, **(IC5)**, **(IC6)**, **(IC6b)**, **(Maj)** and **(Arb)** are not satisfied in general.

**Proof:** For space reasons, we provide only the less obvious proofs.

**(IC2)** Suppose  $k = 0.5$ ,  $d = d_H$  and  $[K_1] = \{00\}$ ,  $[K_2] = \{00, 01\}$  and  $[K_3] = \{00, 01\}$ . We have  $d_m^{d_H, 0.5}(00, \{K_1, K_2, K_3\}) = 0$  and  $d_m^{d_H, 0.5}(01, \{K_1, K_2, K_3\}) = 0$ : **(IC2)** is not satisfied. Note that for  $k \in ]\frac{n-1}{n}, 1]$  and any distance  $d$ , **(IC2)** is satisfied (the aggregation function *max* is retrieved).

**(IC4)** Suppose  $K_1 \models \mu$ ,  $K_2 \models \mu$  and  $\Delta_{\mu}^{d, med^k}(\{K_1, K_2\}) \wedge K_1$  consistent.

Let  $\omega_1 \models \Delta_{\mu}^{d, med^k}(\{K_1, K_2\}) \wedge K_1$ . Then  $d_m^{d,k}(\omega_1, \{K_1, K_2\}) = med^k(d(\omega_1, K_1), d(\omega_1, K_2)) = med^k(0, d(\omega_1, K_2)) = med^k(0, d(\omega_1, \omega_2))$ , where  $\omega_2 \models K_2$  s.t.  $d(\omega_1, K_2) = d(\omega_1, \omega_2)$ . Then  $d(\omega_2, K_1) \leq d(\omega_1, \omega_2)$ :  $d_m^{d,k}(\omega_2, \{K_1, K_2\}) = med^k(d(\omega_2, K_1), 0) \leq d_m^{d,k}(\omega_1, \{K_1, K_2\})$ . As a consequence  $\omega_2$  is selected.

**(IC5)** and **(IC6)** Suppose  $k = 0.5$ , and  $\omega_1$  s.t.  $d_m^{d, 0.5}(\omega_1, E_1) = med^{0.5}(0, 0) = 0$ ,  $d_m^{d, 0.5}(\omega_1, E_2) = med^{0.5}(3, 4, 4) = 4$  and accordingly  $d_m^{d, 0.5}(\omega_1, E_1 \sqcup E_2) = med^{0.5}(0, 0, 3, 4, 4) = 3$ . Suppose  $\omega_2$  s.t.  $d_m^{d, 0.5}(\omega_2, E_1) = med^{0.5}(1, 1) = 1$ ,  $d_m^{d, 0.5}(\omega_2, E_2) = med^{0.5}(2, 6, 7) = 6$  and  $d_m^{d, 0.5}(\omega_2, E_1 \sqcup E_2) = med^{0.5}(1, 1, 2, 6, 7) = 2$ . Then with  $[\mu] = \{\omega_1, \omega_2\}$ ,  $[\Delta_{\mu}^{d, med^{0.5}}(E_1)] = \{\omega_1\}$  and  $[\Delta_{\mu}^{d, med^{0.5}}(E_2)] = \{\omega_1\}$  (and then  $\Delta_{\mu}^{d, med^{0.5}}(E_1)$  and  $\Delta_{\mu}^{d, med^{0.5}}(E_2)$  are jointly consistent), whereas  $[\Delta_{\mu}^{d, med^{0.5}}(E_1 \sqcup E_2)] = \{\omega_2\}$ . This example shows that neither **(IC5)** nor **(IC6)** is satisfied.

**(Arb)** Suppose  $0 < k < 0.5$  (so that  $\Delta^{d, med^k}$  is equivalent to  $\Delta^{d, min}$ ). The following example shows that **(Arb)** is not satisfied. Let  $\omega_1, \omega_2, \omega_3$  be three interpretations s.t.  $d(\omega_1, \omega_3) = 1$ ,  $d(\omega_1, \omega_2) = 1$  and  $d(\omega_2, \omega_3) = 2$ . The bases  $K_1$  and  $K_2$  are defined by  $K_1 = \{\omega_2\}$  and  $K_2 = \{\omega_3\}$ ; and the constraints  $\mu_1$  and  $\mu_2$  are defined by  $\mu_1 = \{\omega_1, \omega_3\}$ ,  $\mu_2 = \{\omega_1, \omega_2\}$ . Then we have:

$$\left. \begin{aligned} [\Delta_{\mu_1}(K_1)] &= [\Delta_{\mu_2}(K_2)] = \{\omega_1\} \\ [\Delta_{\mu_1 \leftrightarrow \neg \mu_2}(\{K_1, K_2\})] &= [\mu_1 \leftrightarrow \neg \mu_2] = \{\omega_2, \omega_3\} \\ \mu_1 \not\models \mu_2 \text{ and } \mu_1 \not\models \mu_2 & \end{aligned} \right\}$$

but  $[\Delta_{\mu_1 \vee \mu_2}(\{K_1, K_2\})] = \{\omega_2, \omega_3\}$  and  $[\Delta_{\mu_1}(K_1)] = \{\omega_1\}$ : contradiction.  $\square$

It turns out that  $k$ -median operators satisfy some but not all the expected rationality properties. In particular the very natural postulate **(IC2)**, asking that the result of the merging is just the conjunction of the bases when this conjunction is consistent, is not satisfied (except for values of  $k$  making the  $k$ -median identical to *max*). We also have:

**Proposition 10** If  $k \geq 0.5$ , then  $\Delta^{d, med^k}$  satisfies **(Arb)**.

No other equity condition is satisfied by such operators:

**Proposition 11** Whatever  $k$ ,  $\Delta^{d, med^k}$  does not satisfy **(PD)** or **(SHE)**.

**Proof:** Consider the following counter-example, for  $k < 0.5$ :  $med^k(1, 3) = 1$  and  $med^k(2, 2) = 2$ :  $(1, 3) <_{med^k} (2, 2)$ : none of **(PD)** or **(SHE)** is satisfied.

Consider the following counter-example, for  $0.5 \leq k < 1$ :  $med^k(0, 4, 4, 7) = 4$  and  $med^k(0, 4, 5, 6) = 5$   $(0, 4, 4, 7) <_{med^k} (0, 4, 5, 6)$ : none of **(PD)** or **(SHE)** is satisfied.

Consider the following counter-example, for  $k = 1$ :  $med^k(0, 4, 4, 7) = 7$  and  $med^k(0, 3, 4, 8) = 8$

$(0, 4, 4, 7) <_{med^k} (0, 3, 4, 8)$ : none of **(PD)** or **(SHE)** is satisfied.  $\square$

Let us now turn to the  $k$ -leximedian operators, that satisfy more expected properties:

**Definition 18** Let  $E = \{K_1, \dots, K_n\}$  be a profile,  $\mu$  an integrity constraint,  $d$  a distance between interpretations and  $k \in ]0, 1[$  a real number. Define  $d_{leximed}^k(\omega, E) = leximed^k(d(\omega, K_1), d(\omega, K_2), \dots, d(\omega, K_n))$ . Then

$$[\Delta_{\mu}^{d, leximed^k}(E)] = \{\omega \models \mu \mid d_{leximed^k}(\omega, E) \text{ is minimal}\}^7$$

Again, some standard operators are recovered by considering specific values of  $k$ :  $\Delta^{d, leximed^k}$  with  $k \in ]0, \frac{1}{n}[$  corresponds to the *leximin* operator  $\Delta^{d, leximin}$ , and  $\Delta^{d, leximed^1}$  to the *leximax* operator  $\Delta^{d, leximax}$ .

As expected  $\Delta^{d, leximed^k}$  operators satisfy more interesting properties than  $\Delta^{d, med^k}$  ones:

**Proposition 12** For any distance  $d$  and any  $k \in ]0, 1[$ ,  $\Delta^{d, leximed^k}$  is a pre-IC merging operator.

We do not have better than that. In particular, these operators are not IC merging ones in the general case:

**Proposition 13**  $\Delta^{d, leximed^k}$  does not satisfy any of **(IC5)**, **(IC6)**, **(Maj)** and **(Arb)** in general.

Concerning egalitarian properties, we reach the arbitration property, but not the other ones:

**Proposition 14** If  $k \geq 0.5$ , then  $\Delta^{d, leximed^k}$  satisfies **(Arb)** but does not satisfy **(PD)** or **(SHE)** in general.

It is easy to explain why the condition  $k \geq 0.5$  is needed. Indeed, we know that when  $k$  goes towards 0 the *leximed<sup>k</sup>* function goes towards *leximin*, and that when  $k$  goes towards 1 *leximed<sup>k</sup>* goes towards *leximax*. So it is natural to obtain an egalitarian behaviour for values between ‘‘classical’’ median ( $k = 0.5$ ) and *leximax*.

Considering Proposition 7, Proposition 14 shows that relaxing some IC postulates is a way for escaping from the *leximax*-based operators while satisfying some egalitarian condition.

## Cumulative Sum Merging Operators

A very convenient representation of the inequalities of a distribution of income is the Lorenz curve (Lorenz 1905). The principle is to focus on the poorest (least satisfied) agents, by looking first at the utility of the poorest one, then at the sum of the utilities of the two poorest ones, etc. To be more precise, on a Lorenz curve, each element  $k$  of the  $x$ -axis corresponds to the  $k$  poorest agents and the value associated with it on the  $y$ -axis is the sum of the utilities of those agents.

This curve can be interpreted in different ways to measure how much a distribution is fair. In particular, the fairest distribution is when for any  $n$ , the  $n\%$  poorest agents own  $n\%$  of the income. The well-known Gini coefficient (Gini 1921; Sen 1973; Dutta 2002), one of the main inequality measures,

<sup>7</sup>w.r.t. the lexicographic order.

	$K_1$	$K_2$	$K_3$	Cum. Sat.	$\Sigma$
000	3	2	2	(2, 4, 7)	13
001	2	3	1	(1, 3, 6)	10
010	2	1	1	(1, 2, 4)	7
011	1	2	0	(0, 1, 3)	4
100	3	1	3	(1, 4, 7)	12
101	2	2	2	(2, 4, 6)	12
110	2	0	2	(0, 2, 4)	6
111	1	1	1	(1, 2, 3)	6

Table 2: Merging with  $\Delta^{CS(d_H, \Sigma)}$

is the (double of the) area between the Lorenz curve and the fairest distribution.

In the following, we adhere to the notion of cumulative sum for defining a new family of merging operators. We translate the distance between a base and an interpretation into a satisfaction value by reversing the scale. Then we compute the cumulative satisfaction vector and take advantage of an aggregation function on it.

Let us define formally the cumulative sum merging operators:

**Definition 19** Let  $d$  be a distance between interpretations,  $f$  an aggregation function,  $E$  a profile and  $\mu$  an integrity constraint. Let  $M = \max(\{d(\omega, \omega') \mid \omega, \omega' \in \Omega\})$ . For an interpretation  $\omega$ , we consider the vector  $(l_{\sigma(1)}, \dots, l_{\sigma(n)})$  where  $l_i = M - d(\omega, K_i)$  is the satisfaction value of agent  $i$  for the interpretation  $\omega$ , and  $\sigma$  is the permutation of  $\{1, \dots, n\}$  sorting the  $l_i$  in ascending order (the less the least satisfied). Then we define the vector of cumulated satisfaction of  $\mu$ ,  $W_d(\omega, E) = (W_1, \dots, W_n)$ , where  $W_i = \sum_{k=1}^i l_{\sigma(k)}$ . Finally, the selected interpretations are the ones which maximize the cumulated satisfaction  $W_d(\omega, E)$ :

$$[\Delta_{\mu}^{CS(d, f)}(E)] = \{\omega \models \mu \mid f(W_d(\omega, E)) \text{ is maximal}\}.$$

Let us illustrate the behavior of a cumulative sum merging operator on a simple example:

**Example 3** We step back to Example 2. The computations are presented in Table 2, in which the values represent satisfaction values of the bases (the more the best), and not anymore distances (the less the best).

We get  $[\Delta_{\top}^{CS(d_H, \Sigma)}(E)] = \{000\}$ . The selected interpretation is 000, because the sum of the cumulative satisfaction vector gives the maximal value of 13.

We recover some existing operators as cumulative sum ones:

**Proposition 15** Let  $d$  be any distance,

- $\Delta^{CS(d, leximin)} = \Delta^{d, leximax}$ .
- $\Delta^{CS(d, leximax)} = \Delta^{d, leximin}$ .

Let us finally turn to the logical properties:

**Proposition 16** • If  $d$  is any distance, and  $f$  is an aggregation function satisfying strict non-decreasingness, then  $\Delta^{CS(d, f)}$  is a pre-IC merging operator.

rank :	$w(\omega, K_i) :$	$w(\omega', K_i) :$
1	$x_1$	$x_1$
...	...	...
$k-1$	$x_{k-1}$	$x_{k-1}$
$k$	$x_k$	$x_{k+1}$
$k+1$	$x_{k+1}$	$x_{k+2}$
...	...	...
$k'-1$	$x_{k'-1}$	$x_{k'}$
$k'$	$x_{k'}$	$x'_k = x_k + g$
$k'+1$	$x_{k'+1}$	$x_{k'+1}$
...	...	...
$l'-1$	$x_{l'-1}$	$x_{l'-1}$
$l'$	$x_{l'}$	$x'_l = x_l - g$
$l'+1$	$x_{l'+1}$	$x_{l'}$
$l-1$	$x_{l-1}$	$x_{l-2}$
$l$	$x_l$	$x_{l-1}$
$l+1$	$x_{l+1}$	$x_{l+1}$
...	...	...
$n$	$x_n$	$x_n$

- $\Delta^{CS(d,f)}$  does not satisfy **(Arb)** and **(Maj)** in the general case.
- $\Delta^{CS(d,f)}$  does not satisfy **(IC5)** and **(IC6)** in the general case.

While **(Arb)** is not satisfied in general, the Pigou-Dalton condition is ensured for instance when any sum of the  $m$ -powers,  $m$  varying, is used as aggregation function:

**Proposition 17** For any integer  $m$ ,  $\Delta^{CS(d,\Sigma^m)}$  satisfies **(PD)**, but **(SHE)** is not satisfied.

**Proof:**

$\Delta^{CS(d,\Sigma^m)}$  satisfies **(PD)**.

Consider  $\omega$  and  $\omega'$  s.t.  $\exists k$  and  $l$  s.t.  $d(\omega, K_k) < d(\omega', K_k) \leq d(\omega', K_l) < d(\omega, K_l)$  and  $d(\omega', K_k) - d(\omega, K_k) = d(\omega, K_l) - d(\omega', K_l)$  and  $\forall i \neq k$  and  $i \neq l$ ,  $d(\omega, K_i) = d(\omega', K_i)$ . We note  $g$  the value  $d(\omega', K_k) - d(\omega, K_k)$ , so we have  $d(\omega', K_k) = d(\omega, K_k) + g$  and  $d(\omega', K_l) = d(\omega, K_l) - g$ .  $g$  is strictly positive.

We note  $\forall i, x_i = n - d(\omega, K_i)$ ,  $x'_k = n - d(\omega', K_k)$  and  $x'_l = n - d(\omega', K_l)$ . We suppose that the  $x_i$  are ordered in the ascending way:  $x_1 \leq x_2 \leq \dots \leq x_n$ .

Let  $l'$  and  $k'$  be the relative ranks of  $x'_l$  and  $x'_k$  into the  $x_i$ -th. We suppose  $l' \neq l$  and  $k' \neq k$ . As  $x_l > x'_l \geq x'_k > x_k$ , we know that if  $l' \neq l$ ,  $l' < l$  and if  $k' \neq k$ ,  $k' > k$ . We have then the following table, if  $k' \neq k$  and  $l' \neq l$ :

From rank 1 to rank  $k-1$ , the  $x_i$ -th are identical so the  $W_i^m$  also.

From rank  $k$  to rank  $k'-1$ , we have in  $\Sigma^m(W_d(\omega, E))$ :

$$\Sigma_{k \dots (k'-1)}^m(W_d(\omega, E)) = (x_1 + \dots + x_{k-1} + x_k)^m + \dots + (x_1 + \dots + x_{k-1} + x_k + \dots + x_{k'-1})^m$$

and in  $\Sigma^m(W_d(\omega', E))$ :

$$\begin{aligned} \Sigma_{k \dots (k'-1)}^m(W_d(\omega', E)) = & (x_1 + \dots + x_{k-1} + x_{k+1})^m + \dots \\ & + (x_1 + \dots + x_{k-1} + x_{k+1} + \dots + x'_k)^m \end{aligned} \quad (3)$$

In Equation 3, the values from  $x_k$  to  $x_{k'-1}$  are replaced by the values from  $x_{k+1}$  to  $x_{k'}$ . Since the  $x_i$ -th are increasing, we have  $\Sigma_{k \dots (k'-1)}^m(W_d(\omega, E)) \leq \Sigma_{k \dots (k'-1)}^m(W_d(\omega', E))$ .

From rank  $k'$  to rank  $l'-1$ , we have in  $\Sigma^m(W_d(\omega, E))$ :

$$\Sigma_{k' \dots (l'-1)}^m(W_d(\omega, E)) = (x_1 + \dots + x_{k-1} + x_k + \dots + x_{k'})^m + \dots + (x_1 + \dots + x_{k-1} + x_k + \dots + x_{l'-1})^m$$

and in  $\Sigma^m(W_d(\omega', E))$ :

$$\begin{aligned} \Sigma_{k' \dots (l'-1)}^m(W_d(\omega', E)) = & (x_1 + \dots + x_{k-1} + x_{k+1} + \dots \\ & + x_{k'-1} + x'_k)^m + \dots + (x_1 + \dots + x_{k-1} + x_{k+1} + \dots \\ & + x'_k + x_{k'+1} + \dots + x_{l'-1})^m = (x_1 + \dots + x_{k-1} + x_{k+1} + \\ & \dots + x_{k'-1} + x_k + g)^m + \dots + (x_1 + \dots + x_{k-1} + x_{k+1} \\ & + \dots + x_k + g + x_{k'+1} + \dots + x_{l'-1})^m \end{aligned}$$

(because  $x'_k = x_k + g$ ). Each term of  $\Sigma^m(W_d(\omega', E))$  is equal to one term of  $\Sigma^m(W_d(\omega, E))$  plus  $g$ . As  $g > 0$ , we have:  $\Sigma_{k' \dots (l'-1)}^m(W_d(\omega, E)) < \Sigma_{k' \dots (l'-1)}^m(W_d(\omega', E))$ .

From rank  $l'$  to rank  $l$ , we have in  $\Sigma^m(W_d(\omega, E))$ :

$$\begin{aligned} \Sigma_{l' \dots l}^m(W_d(\omega, E)) = & (x_1 + \dots + x_k + x_{k+1} + \dots + x_{l'-1} \\ & + x_{l'})^m + \dots + (x_1 + \dots + x_{l'-1} + x_{l'} + \dots + x_{l-1} + x_l)^m \end{aligned}$$

and in  $\Sigma^m(W_d(\omega', E))$ :

$$\begin{aligned} \Sigma_{l' \dots l}^m(W_d(\omega', E)) = & (x_1 + \dots + x_k + g + x_{k+1} + \dots + x_{l'-1} \\ & + x'_l)^m + \dots + (x_1 + \dots + x_k + g + x_{k+1} + \dots + x'_l + x_{l'} + \\ & \dots + x_{l-2} + x_{l-1})^m = (x_1 + \dots + x_k + g + x_{k+1} + \dots \\ & + x_{l'-1} + x_l - g)^m + \dots + (x_1 + \dots + x_k + g + x_{k+1} + \dots \\ & + x_l - g + x_{l'} + \dots + x_{l-1})^m \end{aligned}$$

because  $x'_l = x_l - g$ , so:

$$\begin{aligned} \Sigma_{l' \dots l}^m(W_d(\omega', E)) = & (x_1 + \dots + x_k + x_{k+1} + \dots + x_{l'-1} \\ & + x_l)^m + \dots + (x_1 + \dots + x_k + x_{k+1} + \dots + x_l + x_{l'} + \dots + x_{l-1})^m \end{aligned}$$

In each term of  $\Sigma_{l' \dots l}^m(W_d(\omega', E))$ , compared with  $\Sigma_{l' \dots l}^m(W_d(\omega, E))$ , one element  $x_{l'+i}$  with  $i$  varying from 0 to  $l-l'$  of the sum  $W_j$  is replaced by  $x_l$ . Since the  $x_i$ -th are increasing,  $x_{l'+i} \leq x_l$  for  $0 \leq i \leq l-l'$ , so  $\Sigma_{l' \dots l}^m(W_d(\omega, E)) \leq \Sigma_{l' \dots l}^m(W_d(\omega', E))$ .

From rank  $l+1$  to rank  $n$ , we have in  $\Sigma^m(W_d(\omega, E))$ :

$$\begin{aligned} \Sigma_{(l+1) \dots n}^m(W_d(\omega, E)) = & (x_1 + \dots + x_{l'-1} + x_{l'} + \dots + x_{l-1} \\ & + x_{l+1})^m + \dots + (x_1 + \dots + x_{l'-1} + x_{l'} + \dots \\ & + x_{l-1} + x_l + x_{l+1} + \dots + x_n)^m \end{aligned}$$

and in  $\Sigma^m(W_d(\omega', E))$ :

$$\Sigma_{(l+1) \dots n}^m(W_d(\omega', E)) = (x_1 + \dots + x_k + x_{k+1} + \dots + x_{l'-1} + x_{l+1})^m + \dots + (x_1 + \dots + x_k + x_{k+1} + \dots + x'_l + x_{l'} + \dots + x_n)^m$$

So  $\Sigma_{(l+1) \dots n}^m(W_d(\omega', E)) = \Sigma_{(l+1) \dots n}^m(W_d(\omega, E))$ .

Finally, we obtain  $\Sigma^m(W_d(\omega, E)) < \Sigma^m(W_d(\omega', E))$ , for all integer  $m$ , and  $\omega'$  is selected : **(PD)** is satisfied.

For the other cases, when  $x'_l$  (resp.  $x'_k$ ) has the same rank as  $x_l$  (resp.  $x_k$ ), there are less cases to be studied, the table is simpler, but the result still holds.  $\square$

## Conclusion

In this paper, we have investigated alternative egalitarian conditions to the arbitration postulate. Especially, we have translated to the belief merging framework two egalitarian conditions: Sen-Hammond equity, and Pigou-Dalton property.

We have shown that the distance-based merging operators satisfying Sen-Hammond equity are mainly those for which *leximax* is the aggregation function. This led us to introduce a new family of belief merging operators, the pre-IC operators, which includes the family of IC merging operators as a specific case. We have pointed out a representation theorem for this family, which allows us to define easily some distance-based pre-IC operators. In order to enrich the family of egalitarian merging operators, we have considered two new families of belief merging operators, based respectively on the median and on an aggregated sum (Lorenz curves). We have shown that the operators based on the *leximed<sup>k</sup>* aggregation functions (with  $k \geq 0.5$ ) are pre-IC operators which satisfy the arbitration postulate (**Arb**). We have also proved that the  $\Delta^{CS(d, \Sigma^n)}$  operators from the cumulative sum family are pre-IC operators satisfying (**PD**).

Besides theory-oriented results, this work produced two interesting families of egalitarian operators: cumulative sum ones and *leximed<sup>k</sup>* ones. Before this work the only known egalitarian merging operators were the *leximax*-based ones. Egalitarian operators are significant for all applications where consensual results are expected, i.e., all agents are supposed to be satisfied in the best way, contrariwise to utilitarian/majority operators. Whereas utilitarian operators can be used when the information sources are sensors or databases, egalitarian operators are particularly important in applications where such sources are agents which are autonomous enough to reject the result of the merging if they consider that it is too far from their position.

## Acknowledgments

This work benefited from the support of the AMANDE project of the French National Research Agency (ANR-13-BS02-0004).

## References

Arrow, K.; Sen, A. K.; and Suzumura, K., eds. 2002. *Handbook of Social Choice and Welfare*, volume 1. North-Holland.

Dalton, H. 1920. The measurement of the inequality of incomes. *Economic Journal* 30:pp. 348–461.

Dutta, B. 2002. Inequality, Poverty and Welfare. In Arrow, K. J.; Sen, A. K.; and Suzumura, K., eds., *Handbook of Social Choice and Welfare*. Elsevier. 597–633.

Everaere, P.; Konieczny, S.; and Marquis, P. 2010a. Disjunctive merging: Quota and gmin merging operators. *Artificial Intelligence* 174(12-13):824–849.

Everaere, P.; Konieczny, S.; and Marquis, P. 2010b. The epistemic view of belief merging: Can we track the truth? In *Proceedings of the Nineteenth European Conference on Artificial Intelligence (ECAI'10)*, 621–626.

Gini, C. 1921. Measurement of inequality of incomes. *The Economic Journal* 31(121):pp. 124–126.

Hammond, P. J. 1976. Equity, Arrow's conditions, and Rawls' difference principle. *Econometrica* 44(4):pp. 793–804.

Harsanyi, J. C. 1955. Cardinal welfare, individualistic ethics, and interpersonal comparisons of utility. *Journal of Political Economy* 63:309.

Konieczny, S., and Pino Pérez, R. 2002a. Merging information under constraints: a logical framework. *Journal of Logic and Computation* 12(5):773–808.

Konieczny, S., and Pino Pérez, R. 2002b. On the frontier between arbitration and majority. In *Proceedings of KR'02*, 109–118.

Konieczny, S.; Lang, J.; and Marquis, P. 2004. DA<sup>2</sup> merging operators. *Artificial Intelligence* 157:49–79.

Lin, J., and Mendelzon, A. O. 1999. Knowledge base merging by majority. In *Dynamic Worlds: From the Frame Problem to Knowledge Management*, volume 12 of *Applied Logic Series*, 195–218. Kluwer.

Lorenz, M. O. 1905. Methods of measuring the concentration of wealth. *Publications of the American Statistical Association* 9(70):pp. 209–219.

Moulin, H. 1988. *Axioms of Cooperative Decision Making*. Econometric society monographs. Cambridge University Press. chapter 9.

Rawls, J. 1971. *A Theory of Justice*. Harvard University Press.

Revesz, P. Z. 1997. On the semantics of arbitration. *International Journal of Algebra and Computation* 7(2):133–160.

Sen, A. 1973. *On Economic Inequality*. Oxford University Press.

Sen, A. 1997. *Choice, Welfare and Measurement*. Harvard University Press.

Sen, A. 2005. Social choice theory. In Arrow, K. J., and Intriligator, M., eds., *Handbook of Mathematical Economics*, volume 3. Elsevier. 1073–1181.

Suzumura, K. 1983. *Rational choice, collective decisions and social welfare*. Cambridge University Press.